

Classification of Incidence Scrolls (II)

Rosa Cid-Muñoz

Manuel Pedreira

Authors' address: Departamento de Algebra, Universidad de Santiago de Compostela.
15782 Santiago de Compostela. Galicia. Spain. Phone: 34-81563100-ext.13152. Fax: 34-81597054. e-mail: rosacid@usc.es; pedreira@zmat.usc.es

Abstract: The aim of this paper is to obtain a classification of the scrolls in $\mathbb{P}^n$ which are defined by a one-dimensional family of lines meeting a certain set of linear spaces in $\mathbb{P}^n$, a first classification for genus 0 and 1 is given in paper [1]. These ruled surfaces are called *incidence scrolls*, and such an indicated set is a *base* of the incidence scroll. In particular, we compute its degree and genus. For this, we define the *fundamental incidence scroll* to be the scroll in $\mathbb{P}^n$ formed by the lines which meet $(2n - 3) \mathbb{P}^{n-2}$'s in general position. Then all the others incidence scrolls will be portions of degenerate forms of this. In this way, we can obtain all the incidence scrolls in $\mathbb{P}^n$, $n \geq 3$, with base in general position.

Mathematics Subject Classifications: Primary, 14J26; secondary, 14H25, 14H45.

Introduction: Throughout this paper, the base field for algebraic varieties is $\mathbb{C}$. Let $\mathbb{P}^n$ be the n -dimensional complex projective space and $G(l, n)$ the Grassmannian of l -planes in $\mathbb{P}^n$. Then $R_g^d \subset \mathbb{P}^n$ denotes a scroll of degree d and genus g . We will follow the notation and terminology of [6].

It is useful to represent a scroll in $\mathbb{P}^n$ by a curve $C \subset G(1, n) \subset \mathbb{P}^N$. The lines which intersect a given subspace $\mathbb{P}^r \subset \mathbb{P}^n$ are represented by the points of the special Schubert variety $\Omega(\mathbb{P}^r, \mathbb{P}^n)$. Each $\Omega(\mathbb{P}^r, \mathbb{P}^n)$ is the intersection of $G(1, n)$ with a certain subspace of $\mathbb{P}^N$. Since $G(1, n)$ has dimension $2n - 2$ and we search a curve, we must impose $2n - 3$ linear conditions on $G(1, n)$. Consequently, the choice of subspaces is not arbitrary. Any set of subspaces of $\mathbb{P}^n$ which imposes $2n - 3$ linear conditions on $G(1, n)$ is the base of a certain incidence scroll. The background about Schubert varieties can be found in [8].

The aim of this paper is to obtain a classification of the scrolls in $\mathbb{P}^n$ which are defined by a one-dimensional family of lines meeting a certain set of subspaces of $\mathbb{P}^n$, a first classification for genus 0 and 1 is given in paper [1]. These ruled surfaces are called *incidence scrolls*, and such an indicated set is a *base* of the incidence scroll. Unless otherwise stated, we assume that the base spaces are in general position.

For the convenience of the reader we repeat the relevant material from [1] without proof, thus making our exposition self-contained. Accordingly, Section 1 can be viewed as a brief summary of notation, fundamental definitions and technical results which are used for our analysis. Our first step will be to summarize some general properties of ruled surfaces and, in particular, of incidence scrolls. Having revised the notion of incidence scroll, we have compiled

some basic properties of such a scroll, the detailed proofs appear in [1]. We will expose a method to know when a scroll is determined by incidence. The affirmative solution would allow us to obtain a base for each incidence scroll (there is really only one way to choose this base). This is possible because the families of directrix curves provide a natural and intrinsic characterization of the incidence scrolls. The degree of the scroll given by a general base is provided by Giambelli's formula which appears in [5] or by Young tableaux ([4]). Moreover, the study of deformations of a given incidence scroll is a powerful tool in order to simplify our proofs. If the incidence scroll $R_g^d \subset \mathbb{P}^n$ breaks up into $R_{g_1}^{d_1} \subset \mathbb{P}^r$ and $R_{g_2}^{d_2} \subset \mathbb{P}^s$ with δ generators in common, then $d = d_1 + d_2$ and $g = g_1 + g_2 + \delta - 1$.

In Section 2 we define the *fundamental incidence scroll* to be the scroll in $\mathbb{P}^n$ formed by the lines which meet $(2n - 3)\mathbb{P}^{n-2}$ in general position. All the other incidence scrolls will be portions of degenerate forms of this. In this way, our main contribution is to establish the degree and genus of the incidence scroll in $\mathbb{P}^n$ with base $\mathcal{B} = \{i_s \mathbb{P}^{n-s-2}, \dots, i_1 \mathbb{P}^{n-3}, (2n - 3 - \sum_{j=1}^s (j + 1)i_j) \mathbb{P}^{n-2}\}$ in general position, for any $i_1, \dots, i_s$ between suitable limits.

Section 3 is devoted to study of decomposable ruled surfaces. The important point to note here is the fact that we find all the projective models of these which are defined by incidence. After having computed its degree and genus, we restrict our analysis to those indecomposable incidence scrolls in $\mathbb{P}^n$ whose two directrix curves of minimum degree meet in a only point. Each of these is projected from a point of a decomposable incidence scroll keeping the incidence and genus.

There are people who, even if not related to this paper, have played a important role. My thanks to Proff. W. Fulton, F. Sottile, I. Vainsencher and J. Kock. Special thanks to Prof. A. Lascoux for useful advices (use Maple 6 and library ACE) and for precise bibliographic references. The results on this paper belong to the Ph.D. thesis of the first author whose advisor is the second one.

1 Incidence Scrolls

A *ruled surface* is a surface X together with a surjective morphism $\pi: X \longrightarrow C$ to a smooth curve C such that the fibre X_y is isomorphic to $\mathbb{P}^1$ for every point $y \in C$, and such that π admits a section. There exists a locally free sheaf $\mathcal{E}$ of rank 2 on C such that $X \cong \mathbb{P}(\mathcal{E})$ over C . Conversely, every such $\mathbb{P}(\mathcal{E})$ is a ruled surface over C .

A *scroll* is a ruled surface embedded in $\mathbb{P}^n$ in such a way that the fibres f have degree 1. If we take a very ample divisor on X , $H \sim aC_o + \mathbf{b}f$, then the embedding $\Phi_{|H|}: X \longrightarrow R_g^d \subset \mathbb{P}^n = \mathbb{P}(H^0(\mathcal{O}_X(H)))$ determines a scroll when $a = 1$. A scroll $R_g^d \subset \mathbb{P}^n$ is said to be an *incidence scroll* if it is generated by the lines which meet a certain set $\mathcal{B}$ of linear spaces in $\mathbb{P}^n$, or equivalently, if the correspondent curve in $G(1, n)$ is an intersection of special Schubert varieties $\Omega(\mathbb{P}^r, \mathbb{P}^n)$, $0 \leq r < n - 1$. Such a set is called a *base* of the incidence scroll and

such a base will be denoted by:

$$\mathcal{B} = \{\mathbb{P}^{n_1}, \mathbb{P}^{n_2}, \dots, \mathbb{P}^{n_r}\}.$$

We will write it simply $\mathcal{B}$ when no confusion can arise, where $n_1 \leq n_2 \leq \dots \leq n_r$.

Therefore, unless otherwise stated, we will work with linear spaces in general position. By general position we will mean that $(\mathbb{P}^{n_1}, \dots, \mathbb{P}^{n_r}) \in \mathcal{W} = G(n_1, n) \times \dots \times G(n_r, n)$ is contained in a nonempty open subset of $\mathcal{W}$. For simplicity of notation, we abbreviate it to base in general position.

Proposition 1.1 ([1], Proposition 2.4) *The intersection $C = \bigcap_{i=1}^r \Omega(\mathbb{P}^{n_i}, \mathbb{P}^n)$ of special Schubert varieties associated to linear spaces $\mathbb{P}^{n_i}$, $i = 1, \dots, r$, in general position is an irreducible curve of $G(1, n)$ if and only if it verifies the following equality*

$$rn - (n_1 + n_2 + \dots + n_r) - r = 2n - 3 \quad (IS)$$

■

Moreover, the incidence scroll generated by a base $\mathcal{B}$ have degree d if and only if we obtain the following equality of Schubert cycles:

$$\Omega(n_1, n) \cdots \Omega(n_r, n) = d \Omega(0, 2).$$

Using Young tableaux, we can also compute the degree of the scroll. Consider all ways of filling a 2 by $n - 1$ rectangle with n_1 1's, n_2 2's, and so on until n_r r 's, and the one $r + 1$ in the lower right corner. The fillings have the restriction that the numbers must be weakly increase across each row, but strictly increase down each column. The number of such objects is the degree that we desire ([10]).

Furthermore, we present one of the three main theorems of the symbolic formalism, known as Schubert calculus, for solving enumerative problems.

Theorem 1.2 (Pieri's formula) *For all sequences of integers $0 \leq a_0 < \dots < a_l \leq n$ and for $h = 0, \dots, n - l$, the following formula holds in the cohomology ring $H^*(G(l, n); \mathbb{Z})$:*

$$\Omega(a_0, \dots, a_l) \Omega(h, n) = \sum \Omega(b_0, \dots, b_l)$$

where the sum ranges over all sequences of integers $b_0 < \dots < b_l$ satisfying $0 \leq b_0 \leq a_0 < b_1 \leq a_1 < \dots < b_l \leq a_l$ and $\sum_{i=0}^l b_i = \sum_{i=0}^l a_i - (n - l - h)$.

Proof. See [8], p. 1073. ■

Finally, let us mention an important property of degeneration of these scrolls.

Proposition 1.3 ([1], Proposition 3.1) *Let $R_g^d \subset \mathbb{P}^n$ be an incidence scroll with base $\mathcal{B}$ in general position. Suppose that $\mathbb{P}^{n_i} \vee \mathbb{P}^{n_j} = \mathbb{P}^{n-1}$ and have in common $\mathbb{P}^m$, $m = n_i + n_j - n + 1$. Then the scroll breaks up into:*

- $R_{g_1}^{d_1} \subset \mathbb{P}^n$ with base $\dot{\mathcal{B}} = \{\mathbb{P}^m, \mathbb{P}^{n_1}, \dots, \widehat{\mathbb{P}^{n_i}}, \dots, \widehat{\mathbb{P}^{n_j}}, \dots, \mathbb{P}^{n_r}\}$ (which is possibly degenerate);
- $R_{g_2}^{d_2} \subset \mathbb{P}^{n-1}$ with base $\ddot{\mathcal{B}} = \{\dots, \mathbb{P}^{n_{i-1}-1}, \mathbb{P}^{n_i}, \mathbb{P}^{n_{i+1}-1}, \dots, \mathbb{P}^{n_{j-1}-1}, \mathbb{P}^{n_j}, \mathbb{P}^{n_{j+1}-1}, \dots\}$

which have $\kappa \geq 1$ generators in common. Then, $d = d_1 + d_2$ and $g = g_1 + g_2 + \kappa - 1$.

Moreover, if $m = 0$, then the incidence scroll breaks up into a plane and an incidence scroll $R_g^{d-1} \subset \mathbb{P}^{n-1}$ with base $\ddot{\mathcal{B}}$ in general position. ■

If $m = 0$, then shall refer to this particular degeneration as *join* $\mathbb{P}^{n_i}$ and $\mathbb{P}^{n_j}$ (i.e., $n_i + n_j = n - 1$) and to the inverse as *separate* $\mathbb{P}^{n_i}$ and $\mathbb{P}^{n_j}$ (i.e., $n_i + n_j = n$).

From now on, we will talk about a decomposable incidence scroll if the corresponding ruled surface $X = \mathbb{P}(\mathcal{E})$ has $\mathcal{E}$ decomposable. If $\mathcal{E}$ is indecomposable, then we will talk of an indecomposable incidence scroll.

Let $X = \mathbb{P}(\mathcal{E})$ be a ruled surface over the curve C of genus g , determined by a decomposable normalized bundle $\mathcal{E} \cong \mathcal{O}_C \oplus \mathcal{O}_C(\mathfrak{e})$ such that $\deg(-\mathfrak{e}) = e \geq 0$. Let $H \sim C_o + \mathfrak{b}f$ be the very ample divisor on X with $m = \deg(\mathfrak{b})$ which gives the immersion of the ruled surface as the scroll $R_g^d \subset \mathbb{P}^n$ such that $d = 2m - e$ and $n = 2(m - g) - e + 1 + i$, being i the speciality of the scroll ($i = h^1(\mathcal{O}_C(\mathfrak{b})) + h^1(\mathcal{O}_C(\mathfrak{b} + \mathfrak{e}))$). Geometrically, X has two disjoint directrix, denoted by C_o and C_1 , such that $C_1 \sim C_o - \mathfrak{e}f$. Moreover, it is easy to check that these satisfy $\phi_{\mathfrak{b}+\mathfrak{e}}: C_o \rightarrow C_g^{m-e} \subset \mathbb{P}^{m-e-g+i_1}$ and $\phi_{\mathfrak{b}}: C_1 \rightarrow C_g^m \subset \mathbb{P}^{m-g+i_2}$, being $i_1 = h^1(\mathcal{O}_C(\mathfrak{b} + \mathfrak{e}))$ and $i_2 = h^1(\mathcal{O}_C(\mathfrak{b}))$. Therefore $i = i_1 + i_2$ and $\mathbb{P}^{m-e-g+i_1} \cap \mathbb{P}^{m-g+i_2} = \emptyset$.

Proposition 1.4 ([1], Proposition 2.5) *If $R_g^d \subset \mathbb{P}^n$ is a decomposable incidence scroll with base in general position, then $\mathbb{P}^{m-e-g+i_1}$ and $\mathbb{P}^{m-g+i_2}$ are base spaces.* ■

Proposition 1.5 ([1], Proposition 2.7) *Let $R_g^{2m-e} \subset \mathbb{P}^{2(m-g)-e+1+i}$ be a decomposable incidence scroll with base in general position.*

- (a) *If $\mathfrak{e} \not\sim 0$, then there are $(e+2-g+h^1(\mathcal{O}_C(-\mathfrak{e}))) \mathbb{P}^{m-g+i_2}$'s in $\mathcal{B}$, whenever possible, i.e., when $m-g+i_2 + (e+2-g+h^1(\mathcal{O}_C(-\mathfrak{e}))) (m-e-g+i_1) \leq 4(m-g) - 2e + 2i - 1$ and the generic curve in $|C_o - \mathfrak{e}f|$ is irreducible.*
- (b) *If $\mathfrak{e} \sim 0$, then there are $3 \mathbb{P}^{m-e-g+i_1}$'s in $\mathcal{B}$.* ■

A first classification of scrolls of genus 0 and 1 is given in paper [1], Theorems 4.2 and 5.1.

Theorem 1.6 *Let X be a ruled surface over the curve C of genus $g = 0, 1$. Let $H \sim C_o + \mathfrak{b}f$ be a very ample divisor on X with $m = \deg(\mathfrak{b})$ and let $\Phi_H: X \rightarrow R_g^{2m-e} \subset \mathbb{P}^{2m-e-2g+1}$ be the closed immersion defined by $|H|$. Then R_g^{2m-e} is an incidence scroll if and only if it satisfies one of the following conditions:*

1. $g = 0$ and $e = 0, 1$;
2. $g = 0$ and $m = e + 1$;
3. $g = 1$, $e = -1$ and $m = 2$;
4. $g = 1$, $\mathfrak{e} \sim 0$ and $m = 4$;
5. X decomposable, $\mathfrak{e} \not\sim 0$, $g = 1$, $0 \leq e \leq 3$ and $m = e + 3$. ■

Proposition 1.7 *Let $X = \mathbb{P}(\mathcal{E})$ be a decomposable ruled surface over the curve C of genus g . Let $H \sim C_0 + \mathfrak{b}f$ be a very ample divisor on X which gives the immersion of X as the scroll $R_g^{2m-e} \subset \mathbb{P}^{2(m-g)-e+i+1}$ such that $m = \deg(\mathfrak{b})$.*

1. $(e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1) = m - g + i_2 - 1 \text{ (}\star\text{) and } h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) \leq 3 \Rightarrow R \text{ is an incidence scroll with } \mathcal{B} = \{\mathbb{P}^{m-e-g+i_1}, (e+2-g+h^1(\mathcal{O}_C(-\mathfrak{e})) \mathbb{P}^{m-g+i_2})\}.$
2. $(e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1) > m - g + i_2 - 1 \Rightarrow R \text{ is not an incidence scroll.}$

Proof. $(\star)$ is equivalent to $(e - 1 - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1) = e + i_2 - i_1 - 1$. In particular, $1 \leq h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) \leq e + i_2 - i_1 + 2$.

- a. For $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = 1$, we have $2(m - g) - e + i = 1$, which contradicts the fact that $2(m - g) - e + i \geq 2$.
- b. For $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = 2$, we have $m - e - g + i_1 = m - g + i_2 = 1$. Then the incidence scroll is the smooth quadric surface in $\mathbb{P}^3$ with base $\mathcal{B} = \{3\mathbb{P}^1\}$.
- c. For $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = 3$, i.e., $i_1 - e = i_2 - 1$, we have $\mathcal{B} = \{\mathbb{P}^{m-g+i_2-1}, 3\mathbb{P}^{m-g+i_2}\}$. According to Theorem 1.6, we know that $\mathcal{B}$ generates a rational incidence scroll. The proof is completed by showing that if $(\star)$ is true for $g \geq 1$, then $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) \neq 3$. To do this, suppose that $g \geq 1$ and $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = 3$. Under the above conditions, we see that $e - 1 = g - h^1(\mathcal{O}_C(-\mathfrak{e}))$. Then can assume that $e \geq 1$, on the contrary ($e = 0$), we find that $h^1(\mathcal{O}_C(-\mathfrak{e}))$ is equal to g or $g - 1$, hence that $e - 1 \neq g - h^1(\mathcal{O}_C(-\mathfrak{e}))$. Since $e \geq 1$, we take a general point P of R such that $P \notin \mathbb{P}^{m-g+i_2-1}$ and $P \notin \mathbb{P}^{m-g+i_2}$ for some $\mathbb{P}^{m-g+i_2} \in \mathcal{B}$. The projection of R_g^{2m-e} from P is a scroll with at least two directrix curves contain in linear spaces of dimension $m - g + i_2 - 1$. Since $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = 3$, there is a one-dimensional family of directrix curves $D \sim C_o - \mathfrak{e}f$ that contain P . These curves have the same degree, i.e., $m - e = m - 1$. More precisely, this means that $g = 0$ because R is a smooth irreducible scroll of degree $2m - 1$ in $\mathbb{P}^{2m}$ (see [6]; Remark 2.19.2).

For $g = 0$, we know that $i_1 = i_2 = 0$, hence we conclude from $(\star)$ that $e = 1$, and finally that R is rational of degree $2m - 1$. It is known that $R_0^{2m-1} \subset \mathbb{P}^{2m}$ is incidence with base $\mathcal{B} = \{\mathbb{P}^{m-1}, 3\mathbb{P}^m\}$ ([1]).

Next, suppose that $(e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1)) > m - g + i_2 - 1$. For a suitable number η such that $1 < \eta < m_1$, we have:

$$\begin{aligned} m - g + i_2 + \eta(m - e - g + i_1) &\leq 4(m - g) - 2e + 2i - 1 \quad (\star\star) \\ m - g + i_2 + (\eta + 1)(m - e - g + i_1) &> 4(m - g) - 2e + 2i - 1. \end{aligned}$$

Analysis similar to that in Proposition 1.5 shows that if R is defined by incidence, then there are $\eta \mathbb{P}^{m-g+i_2}$'s in the base. Then $\mathcal{B} = \{\mathbb{P}^{m-g-e+i_1}, \eta \mathbb{P}^{m-g+i_2}, \mathbb{P}^{n_1}, \dots, \mathbb{P}^{n_r}\}$ in general position with $m - g + i_2 + 1 \leq n_i \leq 2(m - g) - e + i$ is the base of R . If $(\star\star)$ is an inequality, then $n_1 = \dots = n_r = 2(m - g) - e + i$. Since the scroll has at least one directrix curve $C_g^m \subset \mathbb{P}^{m-g+i_2} \notin \mathcal{B}$ which is linearly independent from the other directrix curves of degree m , we can take a generic hyperplane $\mathbb{P}^{2(m-g)-e+i}$ through $\mathbb{P}^{m-g+i_2}$. Then there are $(m - e)$ lines in $\mathbb{P}^{2(m-g)-e+i-\eta}$ which meet $\mathbb{P}^{n_1-\eta-1}, \dots, \mathbb{P}^{n_r-\eta-1}, \mathbb{P}^{m-e-g+i_1-1}$ and $\eta \mathbb{P}^{m-g+i_2-\eta}$, which is impossible. ■

Remark 1.8 (1.) is still true if we replace $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) \leq 3$ by $g \leq 2$. For $g = 0$ (i.e., $i_1 = i_2 = 0$) and $e \geq 2$, we obtain $m - e = 1$ and $h^0(\mathcal{O}_X(C_o - \mathfrak{e}f)) = e + 2$. Under these conditions, we find $R_0^{e+2} \subset \mathbb{P}^{e+3}$ with $\mathcal{B} = \{\mathbb{P}^1, (e+2) \mathbb{P}^{e+1}\}$ (see Theorem 1.6). The case $e = 1$ has been study in the above proof. For $g = 1$, $(\star)$ is true only for $e = 3$ and $n = 6$ (see Theorem 1.6). Finally, for $g(C) = 2$, is trivial because $(\star)$ is impossible for any $e \geq 0$.

Finally, we will study the inequality $(e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1)) < m - g + i_2 - 1$. If R is an incidence scroll, then there are an $\mathbb{P}^{m-e-g+i_1}$ and $(e - g + 2 + h^1(\mathcal{O}_C(-\mathfrak{e}))) \mathbb{P}^{m-g+i_2}$'s in $\mathcal{B}$. But these spaces are not sufficient, so we must study the following directrix curves of the scroll. These are curves of type $C_g^s \subset \mathbb{P}^{s-g+i_s}$ such that $m - g + i_2 + 1 \leq s - g + i_s \leq 2(m - g) - e + i$ which belong to a linear system $|C_o + (\mathfrak{D} - \mathfrak{e})f|$ where $\deg(\mathfrak{D}) = s - m$. We proceed in three steps.

3.a) If $s - g + i_s < 2(m - g) - e + i - 1$, then let $m_2 \geq 1$ be the number of directrix curves which are linearly independent on $|C_o + (\mathfrak{D} - \mathfrak{e})f|$.

3.a.i) $(e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1) + m_2(2(m - g) - e + i - (s - g + i_s))) < m - g + i_2 - 1 \Rightarrow \{R \text{ is an incidence scroll} \Rightarrow \{\mathbb{P}^{m-e-g+i_1}, (e + 2 - g + h^1(\mathcal{O}_C(-\mathfrak{e}))) \mathbb{P}^{m-g+i_2}, m_2 \mathbb{P}^{s-g+i_s}\} \subset \mathcal{B}\} \Rightarrow \text{to be continued.}$

3.a.ii) $(e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1) + m_2(2(m - g) - e + i - (s - g + i_s))) = m - g + i_2 - 1 \Rightarrow \{R \text{ is an incidence scroll} \Rightarrow \mathcal{B} = \{\mathbb{P}^{m-e-g+i_1}, (e + 2 - g + h^1(\mathcal{O}_C(-\mathfrak{e}))) \mathbb{P}^{m-g+i_2}, m_2 \mathbb{P}^{s-g+i_s}\}\}.$

3.a.iii) $(e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1) + m_2(2(m - g) - e + i - (s - g + i_s))) > m - g + i_2 - 1 \Rightarrow \{R \text{ is an incidence scroll} \Leftrightarrow (e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1) + 2(m - g) - e + i - (s - g + i_s) = m - g + i_2 - 1\} \text{ and } \mathcal{B} = \{\mathbb{P}^{m-e-g+i_1}, m_1 \mathbb{P}^{m-g+i_2}, \mathbb{P}^{s-g+i_s}\}.$

3.b) $s - g + i_s = 2(m - g) - e + i - 1 \Rightarrow \{R \text{ is an incidence scroll} \Rightarrow \mathcal{B} = \{\mathbb{P}^{m-e-g+i_1}, (e + 2 - g + h^1(\mathcal{O}_C(-\mathfrak{e}))) \mathbb{P}^{m-g+i_2}, \eta \mathbb{P}^{s-g+i_s}\} \text{ with } \eta = m - g + i_2 - 1 - (e - g + h^1(\mathcal{O}_C(-\mathfrak{e}))(m - e - g + i_1))\}.$

3.c) $s - g + i_s = 2(m - g) - e + i \Rightarrow R$ is not an incidence scroll.

2 Fundamental Incidence Scroll

We define *the fundamental incidence scroll* of $\mathbb{P}^n$ to be the incidence scroll with base $\mathcal{B}_n = \{(2n - 3) \mathbb{P}^{n-2}\}$, i.e., the curve $C = \bigcap_{i=1}^{2n-3} \Omega_i(\mathbb{P}^{n-2}, \mathbb{P}^n) \subset G(1, n)$. Using [7], p. 364, we deduce that it is a scroll of degree $\frac{1}{n-1} \binom{2n-2}{n}$ and all the other incidence scrolls are portions of degenerate forms of this. Moreover, we find that the degree of a directrix curve contain in an $\mathbb{P}^{n-2}$ is $3 \frac{(2n-4)!}{(n-3)!n!}$. We have the quadric surface in $\mathbb{P}^3$, elliptic quintic scroll generates by the lines which meet $5 \mathbb{P}^2$ in $\mathbb{P}^4$ and so over.

Let $G := G(1, n)$ be the Grassmannian of lines in a fixed $\mathbb{P}^n$. The fixed $\mathbb{P}^n$ induces a trivial bundle $\mathcal{O}_G^{n+1}$ on G of rank $n + 1$ and the $\mathbb{P}^1$ subspaces induce an subbundle S of $\mathcal{O}_G^{n+1}$ of rank 2. So on G we have the universal exact sequence

$$0 \longrightarrow S \longrightarrow \mathcal{O}_G^{n+1} \longrightarrow Q \longrightarrow 0$$

where Q is a vector bundle on G of rank $n-1$, called *the universal quotient bundle* on G and S is called *the universal subbundle* on G . This sequence determines a homomorphism $S \longrightarrow \Omega_G^1 \otimes Q$. By duality, it gives a homomorphism $T_G \longrightarrow S^\vee \otimes Q$ which is an isomorphism. It follows easily that

$$0 \longrightarrow S \otimes S^\vee \longrightarrow \mathcal{O}_G^{n+1} \otimes S^\vee \longrightarrow Q \otimes S^\vee \longrightarrow 0,$$

hence that $\wedge^{n+1} T_G = \wedge^{n+1} (\mathcal{O}_G(1)^{n+1}) = \bigotimes_1^{n+1} \mathcal{O}_G(1) = \mathcal{O}_G(n+1)$, and finally that $\mathcal{O}_G(K_G) = \mathcal{O}_G(-n-1)$, where K_G denotes the canonical divisor on G . Let us apply this to C which is the intersection of G with $2n-3$ generic hyperplanes of $\mathbb{P}^N$. By [6]; II, Exercise 8.4, we obtain $\mathcal{O}_C(K_C) \cong \mathcal{O}_C(n-4)$ and hence

$$g_C = \frac{n-4}{2n-2} \binom{2n-2}{n} - 1.$$

Proposition 2.1 *The fundamental incidence scroll of $\mathbb{P}^n$ has degree $d(n) = \frac{1}{2} \binom{2n-2}{n}$ and genus $g(n) = \frac{n-4}{2n-2} \binom{2n-2}{n} - 1$. Moreover, the degree of the minimum directrix curve is $3 \frac{(2n-4)!}{(n-3)!n!}$. ■*

If $n \geq 5$, then we obtain an indecomposable incidence scroll because $\dim(\mathbb{P}^{n-2} \cap \mathbb{P}^{n-2}) = n-4$. If we take two minimum directrix curves $C \subset \mathbb{P}^{n-2}$ and $\tilde{C} \subset \mathbb{P}^{n-2}$, then $C \cdot \tilde{C} = 2(n-3) \frac{(2n-4)!}{(n-2)!n!} = -e$.

Proposition 2.2 *Let $R_{g(n,i)}^{d(n,i)} \subset \mathbb{P}^n$ be an incidence scroll with base $\mathcal{B}(n, i) = \{i \mathbb{P}^{n-3}, (2n-3-2i) \mathbb{P}^{n-2}\}$ in general position. Then:*

$$1. \ d(n, i) = \sum_{k=0}^i (-1)^k \binom{i}{k} d(n-k),$$

$$2. \quad g(n, i) = \sum_{k=0}^i (-1)^k \binom{i}{k} g(n-k) - \sum_{k_1=0}^{i-1} \sum_{k_2=0}^{k_1} (-1)^{k_2} \binom{k_1}{k_2} d'(n-k_2-1, i-k_1-1).$$

Proof. The proof is by induction on $i \geq 0$. By the above proposition, $d(n, 0) = d(n)$ and $g(n, 0) = g(n)$. Suppose that the incidence scroll with base $\mathcal{B}(n, i-1)$ in general position has degree $d(n, i-1)$ and genus $g(n, i-1)$. Since $2n-3-2(i-1) > 2$, we can suppose that $2\mathbb{P}^{n-2}$'s are contained in a hyperplane, then $\mathcal{B}(n, i-1)$ breaks up into $\mathcal{B}(n, i) \subset \mathbb{P}^n$ and $\mathcal{B}(n-1, i-1) \subset \mathbb{P}^{n-1}$ with $d(n-1, i-1)$ generators in common. Whence,

$$\begin{aligned} 1. \quad d(n, i) &= d(n, i-1) - d(n-1, i-1) = \sum_{k=0}^{i-1} (-1)^k \binom{i-1}{k} d(n-k) - \sum_{k=0}^{i-1} (-1)^k \\ &\quad \binom{i-1}{k} d(n-k-1) = \sum_{k_1=0}^i (-1)^k \binom{i}{k} d(n-k), \\ 2. \quad g(n, i) &= g(n, i-1) - g(n-1, i-1) - d(n-1, i-1) + 1 = \sum_{k=0}^i (-1)^k \binom{i}{k} g(n-k) \\ &\quad - \sum_{k_1=0}^{i-1} \sum_{k_2=0}^{k_1} (-1)^{k_2} \binom{k_1}{k_2} [d(n-k_2-1, i-k_1-1) - 1]. \quad \blacksquare \end{aligned}$$

Theorem 2.3 Let $R_{g(n, i_1, \dots, i_s)}^{d(n, i_1, \dots, i_s)} \subset \mathbb{P}^n$ be an incidence scroll with base $\mathcal{B}(n, i_1, \dots, i_s) = \{i_s \mathbb{P}^{n-s-2}, \dots, i_1 \mathbb{P}^{n-3}, (2n-3-\sum_{j=1}^s (j+1)i_j) \mathbb{P}^{n-2}\}$ in general position. Then

$$\begin{aligned} 1. \quad d(n, i_1, \dots, i_s) &= \sum_{k_1=0}^{i_s} \sum_{k_2=0}^{i_s+i_{(s-1)}-k_1} \dots \sum_{k_s=0}^{\sum_{j=1}^s i_j - k_{(s-1)}} (-1)^{\sum_{j=1}^s k_j} \binom{i_s}{k_1} \binom{i_s+i_{(s-1)}-k_1}{k_2} \\ &\quad \dots \binom{\sum_{j=1}^s i_j - k_{(s-1)}}{k_s} d(n - \sum_{j=1}^s k_j), \\ 2. \quad g(n, i_1, \dots, i_s) &= \sum_{k_1=0}^{i_s} \sum_{k_2=0}^{i_s+i_{(s-1)}-k_1} \dots \sum_{k_s=0}^{\sum_{j=1}^s i_j - k_{(s-1)}} (-1)^{\sum_{j=1}^s k_j} \binom{i_s}{k_1} \binom{i_s+i_{(s-1)}-k_1}{k_2} \\ &\quad \dots \binom{\sum_{j=1}^s i_j - k_{(s-1)}}{k_s} g(n - \sum_{j=1}^s k_j) - \sum_{k_1=0}^{i_s-1} \sum_{k_2=0}^{k_1} (-1)^{k_2} \binom{k_1}{k_2} [d(n-k_2-1, i_1, \dots, i_{(s-3)}, \\ &\quad i_{(s-2)} + k_2, i_{(s-1)} + 1 + k_1 - k_2, i_s - k_1 - 1) - 1] - \sum_{\alpha=3}^{s+1} \left[\sum_{k_1=0}^{i_s} \sum_{k_2=0}^{i_s+i_{(s-1)}-k_1} \dots \right. \\ &\quad \sum_{k_{(\alpha-2)}=0}^{\sum_{j=s-\alpha+3}^s i_j - k_{(\alpha-3)}} \sum_{k_{(\alpha-1)}=0}^{\sum_{j=s-\alpha+2}^s i_j - k_{(\alpha-2)} - 1} \sum_{k_{\alpha}=0}^{k_{(\alpha-1)}} (-1)^{\sum_{j \neq \alpha-1} k_j} \binom{i_s}{k_1} \binom{i_s+i_{(s-1)}-k_1}{k_2} \dots \\ &\quad \left. \binom{\sum_{j=s-\alpha+3}^s i_j - k_{(\alpha-3)}}{k_{(\alpha-2)}} \binom{k_{(\alpha-1)}}{k_{\alpha}} \right] [d(n-1 - \sum_{j \neq \alpha-1} k_j, i_1, \dots, i_{(s-\alpha-1)}, i_{(s-\alpha)} + k_{(\alpha-1)}, \\ &\quad i_{(s-\alpha+1)} + 1 + k_{(\alpha-2)} + k_{(\alpha-1)} - k_{\alpha}, \sum_{j=s-\alpha+2}^s i_j - k_{(\alpha-2)} - k_{(\alpha-1)} - 1) - 1]. \end{aligned}$$

Proof. We proceed by induction on $s \geq 1$. The case $s = 1$ is Proposition 2.2. If $s = 2$, we will prove that the incidence scroll with base $\mathcal{B}(n, i_1, i_2) \subset \mathbb{P}^n$ has

$$\begin{aligned} 1. \quad d(n, i_1, i_2) &= \sum_{k_1=0}^{i_2} \sum_{k_2=0}^{i_1+i_2-k_1} (-1)^{k_1+k_2} \binom{i_2}{k_1} \binom{i_1+i_2-k_1}{k_2} d(n-k_1-k_2), \\ 2. \quad g(n, i_1, i_2) &= \sum_{k_1=0}^{i_2} \sum_{k_2=0}^{i_1+i_2-k_1} (-1)^{k_1+k_2} \binom{i_2}{k_1} \binom{i_1+i_2-k_1}{k_2} g(n-k_1-k_2) - \sum_{k_1=0}^{i_2-1} \sum_{k_2=0}^{k_1} \\ &\quad (-1)^{k_2} \binom{k_1}{k_2} [d(n-k_2-1, i_1+1+k_1-k_2, i_2-k_1-1) - 1] - \sum_{k_1=0}^{i_2} \sum_{k_2=0}^{i_1+i_2-k_1-1} \sum_{k_3=0}^{k_2} \\ &\quad (-1)^{k_1+k_3} \binom{i_2}{k_1} \binom{k_2}{k_3} [d(n-k_3-k_1-1, i_1+i_2-1-k_1-k_2) - 1]. \end{aligned}$$

To do this, we apply induction on $i_2 \geq 0$. It is true for $i_2 = 0$, by Proposition 2.2. Supposing the theorem true for $i_2 - 1 \geq 0$, we prove it for any i_2 . Since $(2n - 3 - 2i_1 - 3i_2) \geq 0$, $(2n - 3 - 2i_1 - 3(i_2 - 1)) \geq 3$. Let us suppose for the moment $i_1 \geq 1$. Then we can consider that $\mathbb{P}^{n-3} \vee \mathbb{P}^{n-2} = \mathbb{P}^{n-1}$, for any $\mathbb{P}^{n-3}, \mathbb{P}^{n-2} \in \mathcal{B}(n, i_1, i_2 - 1)$. Then $\mathcal{B}(n, i_1, i_2 - 1)$ degenerates into $\mathcal{B}(n, i_1 - 1, i_2) \subset \mathbb{P}^n$ and $\mathcal{B}(n - 1, i_1 - 1, i_2 - 1) \subset \mathbb{P}^{n-1}$ with $d(n - 1, i_1, i_2 - 1)$ generators in common. Whence, writing $i_1 + 1$ instead i_1 , we find that

$$\begin{aligned} 1. \quad d(n, i_1, i_2) &= d(n, i_1 + 1, i_2 - 1) - d(n - 1, i_1, i_2 - 1) = \sum_{k=0}^{i_2} (-1)^k \binom{i_2}{k} d(n - k, i_1 + i_2 - k), \\ 2. \quad g(n, i_1, i_2) &= g(n, i_1 + 1, i_2 - 1) - g(n - 1, i_1, i_2 - 1) - d(n - 1, i_1 + 1, i_2 - 1) + 1 = \\ &\quad \sum_{k=0}^{i_2} (-1)^k \binom{i_2}{k} g(n - k, i_1 + i_2 - k) - \sum_{k_1=0}^{i_2-1} \sum_{k_2=0}^{k_1} (-1)^{k_2} \binom{k_1}{k_2} [d(n - k_2 - 1, i_1 + 1 + k_1 - k_2, i_2 - k_1 - 1) - 1]. \end{aligned}$$

For $i_1 = 0$, we can suppose that $2\mathbb{P}^{n-2}$'s of $\mathcal{B}(n, 0, i_2 - 1)$ are containing in a hyperplane. Then we obtain $\mathcal{B}(n, 1, i_2 - 1)$ which verifies the above assumptions.

Assume the theorem holds for $s - 1 \geq 2$. For s , we use induction on $i_s \geq 0$. An argument similar to $s = 2$ shows that the incidence scroll with base $\mathcal{B}(n, i_1, \dots, i_s)$ has

$$\begin{aligned} 1. \quad d(n, i_1, \dots, i_s) &= d(n, i_1, \dots, i_{(s-1)} + 1, i_s - 1) - d(n, i_1, \dots, i_{(s-2)} + 1, \\ &\quad i_{(s-1)}, i_s - 1) = \sum_{k=0}^{i_s} (-1)^k \binom{i_s}{k} d(n - k, i_1, \dots, i_{(s-3)}, i_{(s-2)} + k, i_{(s-1)} + i_s - k), \\ 2. \quad g(n, i_1, \dots, i_s) &= g(n, i_1, \dots, i_{(s-1)} + 1, i_s - 1) - g(n, i_1, \dots, i_{(s-2)} + 1, \\ &\quad i_{(s-1)}, i_s - 1) - d(n, i_1, \dots, i_{(s-1)} + 1, i_s - 1) + 1 = \sum_{k=0}^{i_s} (-1)^k \binom{i_s}{k} g(n - k, \\ &\quad i_1, \dots, i_{(s-3)}, i_{(s-2)} + k, i_{(s-1)} + i_s - k) - \sum_{k_1=0}^{i_s-1} \sum_{k_2=0}^{k_1} (-1)^{k_2} \binom{k_1}{k_2} [d(n - k_2 - 1, i_1, \dots, i_{(s-3)}, i_{(s-2)} + k_2, i_{(s-1)} + 1 + k_1 - k_2, i_s - k_1 - 1) - 1]. \quad \blacksquare \end{aligned}$$

Remark 2.4 When we talk about $\mathcal{B}(n, i_1, \dots, i_s)$, we must assume that $i_j \geq 0$ for $1 \leq j \leq s$ and $2n - 3 - \sum_{j=1}^4 (j+1)i_j \geq 0$. Let us mention two important consequences and, in particular, the last extends the results of [2].

Corollary 2.5 *Let $\mathcal{B}$ be defined by $\mathcal{B}(n, i_1, \dots, i_{(r-2)}, 0, \dots, 0, i_{(n-r-2)} = 1)$ in general position such that $2 \leq r \leq n - 2$. Then $\mathcal{B}$ is the base of an incidence scroll of degree $d(n, i_1, \dots, i_{(r-2)}, 0, \dots, 0, i_{(n-r-2)} = 1)$ and genus $g(n, i_1, \dots, i_{(r-2)}, 0, \dots, 0, i_{(n-r-2)} = 1)$. Moreover, these are all indecomposable incidence scrolls of $\mathbb{P}^n$. ■*

Corollary 2.6 $\mathcal{B}_{i_1, \dots, i_{(r-1)}}^n := \mathcal{B}(n, i_1, \dots, i_{(r-1)}, 0, \dots, 0, i_{(n-r-2)} = 1)$ in general position generates the incidence scroll of degree $d_{i_1, \dots, i_{(r-1)}}^n = d(n, i_1, \dots, i_{(r-1)}, 0, \dots, 0, i_{(n-r-2)} = 1)$ and genus $g_{i_1, \dots, i_{(r-1)}}^n = g(n, i_1, \dots, i_{(r-1)}, 0, \dots, 0, i_{(n-r-2)} = 1)$. Furthermore, these are all incidence scrolls with base $\mathbb{P}^r$. ■

3 Decomposable Incidence Scrolls

Let $\pi: X = \mathbb{P}(\mathcal{E}) \rightarrow C$ be a geometrically ruled surface over a curve C of genus g determined by a decomposable normalized bundle $\mathcal{E} \cong \mathcal{O}_C \oplus \mathcal{O}_C(\mathfrak{e})$. Let $H \sim C_o + \mathfrak{b}f$ be a very ample divisor on X with $m = \deg(\mathfrak{b})$ which gives the immersion of X as the scroll $R_g^d \subset \mathbb{P}^n$. Following the above notation, we will divided the study in two case.

Remark 3.1 In the sequel, when $\sum$ has not meaning, i.e., the lower limit is bigger than the higher limit, we adopt the convention that it is 0. Moreover, we will write $d(r, h_1, \dots, h_s), g(r, h_1, \dots, h_s), \mathcal{B}(r, h_1, \dots, h_s)$ and $\Delta(r, h_1, \dots, h_s)$, instead of $d_e(r, h_1, \dots, h_s), g_e(r, h_1, \dots, h_s), \mathcal{B}_e(r, h_1, \dots, h_s)$ and $\Delta_e(r, h_1, \dots, h_s)$ respectively, when no confusion can arise. Finally, for abbreviation, we will write $\Delta'(r, h_1, \dots, h_s) := \Delta(r, h_1, \dots, h_s) - 1$.

3.1 Incidence Scrolls with $\mathfrak{e} \sim 0$

Suppose $\mathfrak{e} \sim 0$. If $R_g^{2m} \subset \mathbb{P}^{2(m-g)+i+1}$ is an incidence scroll, then there are $3\mathbb{P}^{m-g+i_1}$'s in $\mathcal{B}$ which impose $3(m-g+i_1)$ independent conditions on $G(1, 2(m-g+i_1)+1)$. Writing r instead of $(m-g+i_1)$, we may set up a one-to-one correspondence between incidence scrolls $R \subset \mathbb{P}^{2r+1}$ with $\mathfrak{e} \sim 0$ and partitions of $r-1$, for any $r \geq 1$. The *partition* of a positive integer r is, by definition, a sequence of weakly decreasing positive integers $\lambda = (\lambda_1 \geq \dots \geq \lambda_s)$ such that $\sum_{i=0}^s \lambda_i = r$.

Proposition 3.2 *The number of incidence scroll $R \subset \mathbb{P}^{2r+1}$ with $\mathfrak{e} \sim 0$ is exactly the number of partitions of $r-1$.*

Proof Let $\lambda = (\lambda_1 \geq \dots \geq \lambda_s)$ be a partition of $r-1$. For each λ_i , $1 \leq i \leq s$, we take a generic $\mathbb{P}^{2r-\lambda_i} \subset \mathbb{P}^{2r+1}$ as base space. Hence every λ gives rise an incidence scroll of $\mathbb{P}^{2r+1}$ with base $\mathcal{B} = \{3\mathbb{P}^r, \mathbb{P}^{2r-\lambda_1}, \dots, \mathbb{P}^{2r-\lambda_s}\}$ where $\mathfrak{e} \sim 0$. Conversely, to every $\mathcal{B}_\lambda = \{3\mathbb{P}^r, \mathbb{P}^{n_1}, \dots, \mathbb{P}^{n_s}\} \subset \mathbb{P}^{2r+1}$ such that $r+1 \leq n_1$ there corresponds a partition $(2r-n_1 \geq \dots \geq 2r-n_s)$ of $r-1$. ■

Lemma 3.3 *For each partition $(r-h_1 \geq h_1-h_2 \geq \dots \geq h_{(s-1)}-h_s \geq 1, \binom{h_s-1}{\cdot}, 1)$ of $r-1$, we define $\Delta_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s) := \sigma(r)^3 \cdot \sigma(r+h_1-1) \cdot \sigma(2r-h_1+h_2) \cdot \dots \cdot \sigma(2r-h_{(s-1)}+h_s) \cdot \sigma(2r-1)^{(h_s-1)}$ an intersection of Schubert cycles in $\mathbb{P}^{2r+1}$. Then:*

$$\begin{aligned} 1. \Delta_{\mathfrak{e}_{\sim 0}}(r, h) &= \sum_{k=0}^{h-1} \binom{h-1}{k} (r-2k+1); \\ 2. \Delta_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s) &= \sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \sum_{k_3=0}^{h_{(s-2)}-h_{(s-1)}-2k_2} \dots \sum_{k_{(s-1)}=0}^{h_2-h_{(s-1)}-2 \sum_{i=2}^{s-2} k_i} \\ &\quad \binom{h_s-1}{k_1} (h_1-h_s+1-2 \sum_{j=2}^{s-1} k_j) (r-2k_1-h_1+h_s+1), \text{ for } s \geq 2. \end{aligned}$$

Proof. 1. For $s=1$, we apply induction on $h \geq 1$. Then $\Delta(r, 1) = r+1$ because it is the degree of a directrix curve, which is containing in an $\mathbb{P}^{r+1}$, of the incidence scroll $R_0^{2r} \subset \mathbb{P}^{2r+1}$. Suppose that the lemma is true for $h-1 \geq 1$, we prove it for any h . From Pieri's formula, we have

$$\Delta(r, h) = \Delta(r, h-1) + \Delta(r-2, h-1) = \sum_{k=0}^{h-1} \binom{h-1}{k} \Delta(r-2k, 1).$$

The proof of the last equality is by induction on $h \geq 1$, using the fact that $\binom{h-1}{k} = \binom{h-2}{k} + \binom{h-2}{k-1}$.

2. Suppose $s=2$. We will see in Theorem 3.5 that $\Delta(r, h_1, 1) = h_1(r-h_1+2)$ because the incidence scroll which corresponds to the partition $(r-h_1, h_1-1)$ is a nonspecial scroll of degree $2h_1(r-h_1+1)$ and genus $(r-h_1)(h_1-1)$. In general, using Pieri's formula, $\Delta(r, h_1, h_2) = \Delta(r, h_1-1, h_2-1) + \Delta(r-2, h_1-1, h_2-1) = \sum_{k=0}^{h_2-1} \binom{h_2-1}{k} \Delta(r-2k, h_1-h_2+1, 1) = \sum_{k=0}^{h_2-1} \binom{h_2-1}{k} (h_1-h_2+1)(r-2k-h_1+h_2+1)$.

Assume the formula holds for $s-1 \geq 1$; we will prove it for s . We have divided the proof into 2 steps.

$$\begin{aligned} \text{(a) For } h_s = 1, \text{ according to Pieri's formula, } \Delta(r, h_1, \dots, h_{(s-1)}, 1) &= \sum_{k=0}^{h_{(s-1)}-1} \\ &\quad \sigma(r-2k)^3 \sigma(r+h_1-4k-1) \sigma(2r-h_1+h_2-4k) \dots \sigma(2r-h_{(s-3)}+h_{(s-2)}- \\ &\quad 4k) \sigma(2r-h_{(s-2)}-2k+1) \Big|_{2r-4k+1} = \sum_{k=0}^{h_{(s-1)}-1} \Delta(r-2k, h_1-2k, \dots, h_{(s-2)}- \\ &\quad 2k, 1) \stackrel{\text{induction on } s-1}{=} \sum_{k_1=0}^{h_{(s-1)}-1} \sum_{k_2=0}^{h_{(s-2)}-1-2k_1} \dots \sum_{k_{(s-2)}=0}^{h_2-1-2 \sum_{j=1}^{s-3} k_j} (h_1-2 \sum_{j=1}^{s-2} k_j) (r-h_1+ \\ &\quad 2). \end{aligned}$$

2). The notation $|_k$ means that the intersection is in $\mathbb{P}^k$ instead of $\mathbb{P}^{2r+1}$.

- (b) For $h_s \geq 2$, $\Delta(r, h_1, \dots, h_s) = \Delta(r, h_1 - 1, \dots, h_s - 1) + \Delta(r - 2, h_1 - 1, \dots, h_s - 1)$. Then, for any $h_s \geq 1$, $\Delta(r, h_1, \dots, h_s) = \sum_{k=0}^{h_s-1} \binom{h_s-1}{k} \Delta(r - 2k, h_1 - h_s + 1, \dots, h_{(s-1)} - h_s + 1, 1)$. ■

Proposition 3.4 *Each partition $(r - h, 1^{(h-1)}) := (r - h, 1, \dots, 1)$ of $r - 1$ gives rise to an incidence scroll $R_{g_{\mathfrak{e}_{\sim 0}}(r, h)}^{d_{\mathfrak{e}_{\sim 0}}(r, h)} \subset \mathbb{P}^{2r+1}$ with base $\mathcal{B}_{\mathfrak{e}_{\sim 0}}(r, h) = \{3\mathbb{P}^r, \mathbb{P}^{r+h}, (h-1)\mathbb{P}^{2r-1}\}$ in general position, such that*

- $d_{\mathfrak{e}_{\sim 0}}(r, h) = 2 \sum_{k=0}^{h-1} \binom{h-1}{k} (r - 2k),$
- $g_{\mathfrak{e}_{\sim 0}}(r, h) = \sum_{k_1=2}^h \sum_{k_2=0}^{h-k_1} \binom{h-k_1}{k_2} \Delta'_{\mathfrak{e}_{\sim 0}}(r - 2k_2 - 2, k_1 - 1).$

Proof. From Theorem 1.6, it follows that $d(r, 1) = 2r$ and $g(r, 1) = 0$. Suppose now that $h > 1$. If, in $\mathcal{B}(r, h)$, $\mathbb{P}^{r+h} \vee \mathbb{P}^{2r-1} = \mathbb{P}^{2r}$, then R breaks up into $R_{g(r-2, h-1)}^{d(r-2, h-1)} \subset \mathbb{P}^{2r-3}$ with $\mathcal{B}(r-2, h-1) = \{3\mathbb{P}^{r-2}, \mathbb{P}^{r+h-3}, (h-2)\mathbb{P}^{2r-5}\}$ and $R_{g(r, h-1)}^{d(r, h-1)} \subset \mathbb{P}^{2r+1}$ with $\mathcal{B}(r, h-1) = \{3\mathbb{P}^r, \mathbb{P}^{r+h-1}, (h-2)\mathbb{P}^{2r-1}\}$, and $\Delta(r-2, h-1)$ generators in common. Whence, by induction on $h \geq 1$,

1. $d(r, h) = d(r-2, h-1) + d(r, h-1) = \sum_{k=0}^{h-1} \binom{h-1}{k} d(r-2k, 1),$
2. $g(r, h) = g(r-2, h-1) + g(r, h-1) + \Delta'(r-2, h-1) = \sum_{k_1=2}^h \sum_{k_2=0}^{h-k_1} \binom{h-k_1}{k_2} \Delta'(r-2k_2-2, k_1-1).$ ■

Theorem 3.5 *For $s \geq 2$, each partition $(r-h_1, h_1-h_2, \dots, h_{(s-1)}-h_s, 1^{(h_s-1)})$ of $r-1$ gives rise to $R_{g_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s)}^{d_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s)} \subset \mathbb{P}^{2r+1}$ with $\mathcal{B}_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s) = \{3\mathbb{P}^r, \mathbb{P}^{r+h_1}, \mathbb{P}^{2r-h_1+h_2}, \dots, \mathbb{P}^{2r-h_{(s-1)}+h_s}, (h_s-1)\mathbb{P}^{2r-1}\}$ in general position, such that*

- $d_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s) = 2(h_1 - h_2 + 1) \sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \dots \sum_{k_{(s-1)}=0}^{h_2-h_3} \binom{h_s-1}{k_1} (r - h_1 + h_2 - 2 \sum_{i=1}^{s-1} k_i),$
- $g_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s) = (h_1 - h_2) \sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \dots \sum_{k_{(s-1)}=0}^{h_2-h_3} \binom{h_s-1}{k_1} (r - h_1 + h_2 - 1 - 2 \sum_{i=1}^{s-1} k_i) + \sum_{k_1=2}^{h_s} \sum_{k_2=0}^{h_s-k_1} \binom{h_s-k_1}{k_2} \Delta'_{\mathfrak{e}_{\sim 0}}(r - 2k_2 - 2, h_1 - h_s + k_1 - 1, \dots, h_{(s-1)} - h_s + k_1 - 1, k_1 - 1) + \sum_{\alpha=2}^{s-1} \left[\sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \dots \sum_{k_{(s-\alpha)}=0}^{h_{(\alpha+1)}-h_{(\alpha+2)}} \sum_{k_{(s-\alpha+1)}=0}^{h_\alpha-h_{(\alpha+1)}-1} \binom{h_s-1}{k_1} \Delta'_{\mathfrak{e}_{\sim 0}}(r - 2 - 2 \sum_{i=1}^{s-\alpha+1} k_i, h_1 - h_\alpha + 1, \dots, h_{(\alpha-1)} - h_\alpha + 1, 1) \right].$

Proof. Suppose $h_s = 1$. We first compute $s = 2$. Therefore, $d(r, 2, 1) = 4r - 4$ and $g(r, 2, 1) = r - 2$, by [1], Example 3.3. By induction on $h_1 \geq 1$, we have

- $d(r, h, 1) = d(r, 1) + d(r - 2, h - 1, 1) = 2r + 2(h - 1)(r - h) = 2h(r - h + 1);$
- $g(r, h, 1) = g(r, 1) + g(r - 2, h - 1, 1) + r - 2 = (r - 1 - h)(h - 2) + (r - 2) = (r - h)(h - 1).$

Assuming theorem to hold for $s - 1$, we will prove it for s . To this end, we suppose that, in $\mathcal{B}(r, h_1, \dots, h_{(s-1)}, 1)$, $\mathbb{P}^{r+h_1} \vee \mathbb{P}^{2r-h_{(s-1)}+1} = \mathbb{P}^{2r}$. Then we obtain $\mathcal{B}(r, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1) \subset \mathbb{P}^{2r+1}$ and $\mathcal{B}(r - 2, h_1 - 1, \dots, h_{(s-1)} - 1, 1) \subset \mathbb{P}^{2r-3}$ with $\Delta(r - 2, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1)$ generators in common. In particular, using that $\mathcal{B}(r, h_1, \dots, h_{(s-2)}, 1, 1) = \mathcal{B}(r, h_1, \dots, h_{(s-2)}, 1)$, we deduce that

- $d(r, h_1, \dots, h_{(s-1)}, 1) = d(r, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1) + d(r - 2, h_1 - 1, \dots, h_{(s-1)} - 1, 1) = \sum_{k=0}^{h_{(s-1)}-1} d(r - 2k, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1) = 2(h_1 - h_2 + 1) \sum_{k_1=0}^{h_{(s-1)}-1} \sum_{k_2=0}^{h_{(s-2)}-h_{(s-1)}} \dots \sum_{k_{(s-2)}=0}^{h_2-h_3} (r - h_1 + h_2 - 2 \sum_{i=1}^{s-2} k_i),$
- $g(r, h_1, \dots, h_{(s-1)}, 1) = g(r, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1) + g(r - 2, h_1 - 1, \dots, h_{(s-1)} - 1, 1) + \Delta'(r - 2, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1) = \sum_{k=0}^{h_{(s-1)}-1} g(r - 2k, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1) + \sum_{k=0}^{h_{(s-1)}-2} \Delta'(r - 2(k+1), h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1).$

Finally, we take $h_s \geq 2$. In $\mathcal{B}(r, h_1, \dots, h_s)$, we set that $\mathbb{P}^{r+h_1} \vee \mathbb{P}^{2r-1} = \mathbb{P}^{2r}$. Then $\mathcal{B}(r, h_1, \dots, h_s)$ breaks up into $\mathcal{B}(r, h_1 - 1, \dots, h_s - 1) \subset \mathbb{P}^{2r+1}$ and $\mathcal{B}(r - 2, h_1 - 1, \dots, h_s - 1) \subset \mathbb{P}^{2r-3}$ with $\Delta(r - 2, h_1 - 1, \dots, h_s - 1)$ generators in common. By induction on $h_s \geq 1$, we conclude that

- $d(r, h_1, \dots, h_s) = \sum_{k=0}^{h_s-1} \binom{h_s-1}{k} d(r - 2k, h_1 - h_s + 1, \dots, h_{(s-1)} - h_s + 1, 1),$
- $g(r, h_1, \dots, h_s) = \sum_{k=0}^{h_s-1} \binom{h_s-1}{k} g(r - 2k, h_1 - h_s + 1, \dots, h_{(s-1)} - h_s + 1, 1) + \sum_{k_1=2}^{h_s} \sum_{k_2=0}^{h_s-k_1} \binom{h_s-k_1}{k_2} \Delta'(r - 2(k_2+1), h_1 - h_s + k_1 - 1, \dots, h_{(s-1)} - h_s + k_1 - 1, k_1 - 1).$

■

3.2 Incidence Scrolls with $\mathfrak{e} \not\sim 0$

We can continue with a similar method for provide all the incidence scrolls which have $\mathfrak{e} \not\sim 0$ but $\deg(\mathfrak{e}) = 0$. Writing r instead of $m - g + i_1$, there is a one-to-one correspondence between incidence scrolls in $\mathbb{P}^{2r+1}$ with base $\mathcal{B} = \{2\mathbb{P}^r, \mathbb{P}^{n_1}, \dots, \mathbb{P}^{n_s}\}$ such that $r < n_1$ and partitions of $2r - 1$ such that $\lambda_1 \leq r - 1$.

Lemma 3.6 *Let $\{h_1, \dots, h_s\}$ be a finite number of positive integers such that $1 \leq h_1 \leq r$, $1 \leq h_2 \leq r + h_1$, $1 \leq h_i \leq h_{(i-1)} - 1$ for $3 \leq i \leq s$ and $h_s + \sum_{i=3}^s (h_{(i-1)} - h_i) = h_2$. Suppose $\Delta_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s) :=$*

$$\begin{cases} \sigma(r)^2 \cdot \sigma(r + h_1 - 1) \cdot \sigma(2r - 1)^{(r+h_1-1)}, & s = 1 \\ \sigma(r)^2 \cdot \sigma(r + h_1 - 1) \cdot \sigma(r - h_1 + h_2) \cdot \sigma(2r - h_2 + h_3) \cdot \dots \cdot \sigma(2r - h_{(s-1)} + h_s) \cdot \sigma(2r - 1)^{(h_s-1)}, & s \geq 2 \end{cases}$$

an intersection of Schubert cycles in $\mathbb{P}^{2r+1}$. Then:

$$\begin{aligned} 1. \Delta_{\mathfrak{e}_{\sim 0}}(r, h) &= \begin{cases} d_{\mathfrak{e}_{\sim 0}}(r, r-1), & h = 1 \\ \sum_{k_1=h-1}^r \sum_{k_2=h-2}^{k_1} \dots \sum_{k_{(h-1)}=1}^{k_{(h-2)}} d_{\mathfrak{e}_{\sim 0}}(k_{(h-1)}, k_{(h-1)} - 1), & h > 1 \end{cases} \\ 2. \Delta_{\mathfrak{e}_{\sim 0}}(r, h_1, h_2) &= \sum_{k=0}^{h_2-1} \binom{h_2-1}{k} (r - h_1 + h_2 + 1 - 2k); \\ 3. \Delta_{\mathfrak{e}_{\sim 0}}(r, h_1, \dots, h_s) &= \sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \sum_{k_3=0}^{h_{(s-2)}-h_{(s-1)}-2k_2} \dots \sum_{k_{(s-1)}=0}^{h_2-h_s-2 \sum_{i=2}^{s-2} k_i} \binom{h_s-1}{k_1} \\ &\quad (r - h_1 + h_s + 1 - 2k_1), \text{ for } s \geq 3. \end{aligned}$$

Proof. We only give the main ideas of the proof because it is similar to that of Lemma 3.3.

$$\begin{aligned} 1. \Delta(r, 1) &= d_{\mathfrak{e}_{\sim 0}}(r, r-1) \text{ and } \Delta(r, h) = \Delta(r-1, h) + \Delta(r, h-1) = \sum_{k=h-1}^r \Delta(k, h-1) \\ &\quad (\text{use the fact that } \Delta(r, r+1) = \Delta(r, r)). \\ 2. \text{ For } s = 2, \text{ we have } \Delta(r, h_1, 1) &= \sigma(r - h_1 + 1)^4_{|2(r-h_1+1)+1} = r - h_1 + 2 \text{ and} \\ \Delta(r, h_1, h_2) &= \Delta(r, h_1 - 1, h_2 - 1) + \Delta(r - 1, h_1, h_2 - 1) = \sum_{k_1=0}^{h_2-1} \binom{h_2-1}{k_1} \Delta(r - k, h_1 - h_2 + 1 + k, 1). \\ 3. \text{ When } s \geq 3 \text{ and } h_s = 1, \text{ we obtain } \Delta(r, h_1, \dots, h_{(s-1)}, 1) &= \sum_{k=0}^{h_{(s-1)}-1} \sigma(r-k)^2 \sigma(r+h_1-1-2k) \sigma(r-h_1+h_2-2k) \sigma(2r-h_2+h_3-2k) \dots \sigma(2r - \\ &\quad h_{(s-3)} + h_{(s-2)} - 2k)_{|2(r-k)+1} = \sum_{k=0}^{h_{(s-1)}-1} \Delta(r-k, h_1-k, h_2-2k, \dots, h_{(s-2)} - \\ &\quad 2k, 1) = \sum_{k_1=0}^{h_{(s-1)}-1} \sum_{k_2=0}^{h_{(s-2)}-1-2k_1} \dots \sum_{k_{(s-2)}=0}^{h_2-1-2 \sum_{i=2}^{s-3} k_i} (r - h_1 + 2). \end{aligned}$$

Finally, when $h_s \geq 1$, Pieri's formula makes it obvious that $\Delta(r, h_1, \dots, h_s)$

$$= \sum_{k=0}^{h_s-1} \binom{h_s-1}{k} \Delta(r-k, h_1-h_s+1+k, h_2-h_s+1, \dots, h_{(s-1)}-h_s+1, 1). \quad \blacksquare$$

Proposition 3.7 *Each partition $(r-h, 1^{(r+h-1)})$ of $2r-1$ such that $h \geq 1$ gives rise to $R_{g_{\mathbf{e}_{\sim 0}(r,h)}}^{\mathbf{d}_{\mathbf{e}_{\sim 0}(r,h)}} \subset \mathbb{P}^{2r+1}$ with base $\mathcal{B}_{\mathbf{e}_{\sim 0}(r,h)} = \{2\mathbb{P}^r, \mathbb{P}^{r+h}, (r+h-1)\mathbb{P}^{2r-1}\}$ in general position, where:*

$$\begin{aligned} \bullet \quad d_{\mathbf{e}_{\sim 0}(r,h)} &= \sum_{k_1=h}^r \sum_{k_2=h-1}^{k_1} \cdots \sum_{k_{(h-1)=2}}^{k_{(h-2)}} \sum_{k_h=1}^{k_{(h-1)}} d_{\mathbf{e}_{\sim 0}(k_h, k_h-1), \\ \bullet \quad g_{\mathbf{e}_{\sim 0}(r,h)} &= \sum_{k_1=h}^r \sum_{k_2=h-1}^{k_1} \cdots \sum_{k_{(h-1)=2}}^{k_{(h-2)}} \sum_{k_h=1}^{k_{(h-1)}} g_{\mathbf{e}_{\sim 0}(k_h, k_h-1)} + \sum_{\alpha=2}^h \left[\sum_{k_1=h}^r \right. \\ &\quad \left. \sum_{k_2=h-1}^{k_1} \cdots \sum_{k_{(\alpha-1)=h-\alpha+2}}^{k_{(\alpha-2)}} \sum_{k_\alpha=h-\alpha+1}^{k_{(\alpha-1)}-1} \Delta'_{\mathbf{e}_{\sim 0}}(k_\alpha, h-\alpha+1) \right] + \sum_{k=h}^{r-1} \Delta'_{\mathbf{e}_{\sim 0}}(k, h). \end{aligned}$$

Proof. We proceed by induction on $h \geq 1$. For $h=1$, suppose that, in $\mathcal{B}(r, 1)$, $\mathbb{P}^{r+1} \vee \mathbb{P}^{2r-1} = \mathbb{P}^{2r}$. Then $\mathcal{B}(r, 1)$ degenerates into $\mathcal{B}_{\mathbf{e}_{\sim 0}}(r, r-1) \subset \mathbb{P}^{2r+1}$ and $\mathcal{B}_{\mathbf{e}_{\sim 0}}(r-1, 1) \subset \mathbb{P}^{2r-1}$ with $\Delta_{\mathbf{e}_{\sim 0}}(r-1, 1)$ generators in common. Whence, by induction on $r \geq 2$, we find:

$$\begin{aligned} \bullet \quad d(2, 1) &= d(2, 2) = d_{\mathbf{e}_{\sim 0}}(2, 1) + d_{\mathbf{e}_{\sim 0}}(1, 0) \text{ and } d(r, 1) = d_{\mathbf{e}_{\sim 0}}(r, r-1) + \\ &\quad d(r-1, 1) = \sum_{k=1}^r d_{\mathbf{e}_{\sim 0}}(k, k-1), \\ \bullet \quad g(2, 1) &= g_{\mathbf{e}_{\sim 0}}(2, 1) + g_{\mathbf{e}_{\sim 0}}(1, 0) + \Delta'(1, 1) \text{ and } g(r, 1) = g_{\mathbf{e}_{\sim 0}}(r, r-1) + \\ &\quad g(r-1, 1) + \Delta'(r-1, 1) = \sum_{k=1}^r g_{\mathbf{e}_{\sim 0}}(k, k-1) + \sum_{k=1}^{r-1} \Delta'(k, 1). \end{aligned}$$

Assume the formulas hold for $h-1$; we will prove them for $h > 1$. If $\mathbb{P}^{r+h} \vee \mathbb{P}^{2r-1} = \mathbb{P}^{2r}$, then $\mathcal{B}(r, h)$ breaks up into $\mathcal{B}(r, h-1) \subset \mathbb{P}^{2r+1}$ and $\mathcal{B}(r-1, h) \subset \mathbb{P}^{2r-1}$ with $\Delta_{\mathbf{e}_{\sim 0}}(r-1, h)$ generators in common. Therefore,

$$\begin{aligned} \bullet \quad d(r, h) &= d(r, h-1) + d(r-1, h) = \sum_{k=h}^r d(k, h-1), \\ \bullet \quad g(r, h) &= g(r, h-1) + g(r-1, h) + \Delta'(r-1, h) = \sum_{k=h}^r g(k, h-1) + \sum_{k=h}^{r-1} \Delta'(k, h). \end{aligned}$$

By induction on $r \geq h$, since $\mathcal{B}(r, r) = \mathcal{B}(r, r-1)$, the last equalities are obvious. Combining the two results, we can rewrite:

$$\begin{aligned} \bullet \quad d(r, h) &= \sum_{k_1=h}^r \sum_{k_2=h-1}^{k_1} \cdots \sum_{k_{(h-1)=2}}^{k_{(h-2)}} d(k_{(h-1)}, 1), \\ \bullet \quad g(r, h) &= \sum_{k_1=h}^r \sum_{k_2=h-1}^{k_1} \cdots \sum_{k_{(h-1)=2}}^{k_{(h-2)}} g(k_{(h-1)}, 1) + \sum_{k=h}^{r-1} \Delta'(k, h) + \sum_{\alpha=2}^{h-1} \left[\sum_{k_1=h}^r \right. \\ &\quad \left. \sum_{k_2=h-1}^{k_1} \cdots \sum_{k_{(\alpha-1)=h-\alpha+2}}^{k_{(\alpha-2)}} \sum_{k_\alpha=h-\alpha+1}^{k_{(\alpha-1)}-1} \Delta'(k_\alpha, h-\alpha+1) \right]. \end{aligned}$$

Theorem 3.8 For $s \geq 2$, each partition $(r-h_1, r+h_1-h_2, h_2-h_3, \dots, h_{(s-1)}-h_s, 1^{(h_s-1)})$ of $2r-1$ such that $h_1 \geq 1$ gives rise to the scroll $R_{g\mathfrak{e}_{\neq 0}^{(r, h_1, \dots, h_s)}}^{d\mathfrak{e}_{\neq 0}^{(r, h_1, \dots, h_s)}}$ $\subset \mathbb{P}^{2r+1}$ with base $\mathcal{B}_{\mathfrak{e}_{\neq 0}}(r, h_1, \dots, h_s) = \{2\mathbb{P}^r, \mathbb{P}^{r+h_1}, \mathbb{P}^{r-h_1+h_2}, \mathbb{P}^{2r-h_2+h_3}, \dots, \mathbb{P}^{2r-h_{(s-1)}+h_s}, (h_s-1)\mathbb{P}^{2r-1}\}$ in general position, where

$$\begin{aligned} \bullet \quad d\mathfrak{e}_{\neq 0}^{(r, h_1, \dots, h_s)} &= 2 \sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \dots \sum_{k_{(s-1)}=0}^{h_2-h_3} \binom{h_s-1}{k_1} (r-h_1+h_2-2 \sum_{i=1}^{s-1} k_i), \\ \bullet \quad g\mathfrak{e}_{\neq 0}^{(r, h_1, \dots, h_s)} &= \sum_{\alpha=1}^{s-2} \left[\sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \dots \sum_{k_{\alpha}=0}^{h_{(s-\alpha+1)}-h_{(s-\alpha+2)}} \sum_{k_{(\alpha+1)}=0}^{h_{(s-\alpha)}-h_{(s-\alpha+1)}} \right. \\ &\quad \left. \binom{h_s-1}{k_1} \delta'(r-1 - \sum_{i=1}^{\alpha+1} k_i, h_1-h_{(s-\alpha)}+1 + \sum_{i=1}^{\alpha+1} k_i, h_2-h_{(s-\alpha)}+1, \dots, h_{(s-\alpha-1)}-h_{(s-\alpha)} \right. \\ &\quad \left. + 1, 1) \right] + \sum_{k_1=0}^{h_s-2} \sum_{k_2=0}^{k_1} \binom{k_1}{k_2} \Delta'_{\mathfrak{e}_{\neq 0}}(r-k_2-1, h_1-k_1+k_2, h_2-1-k_1, \dots, h_s-1-k_1); \end{aligned}$$

and, for $s \geq 3$, $\delta(r, h_1, \dots, h_{s-2}, 1) = \sum_{k_1=0}^{h_{(s-2)}-1} \sum_{k_2=0}^{h_{(s-3)}-1-2k_1} \dots \sum_{k_{(s-3)}=0}^{h_2-1-2 \sum_{i=1}^{s-4} k_i} (r-h_1+1) = \delta'(r, h_1, \dots, h_{s-2}, 1) + 1$.

Proof. First, we will compute the case $s = 2$. For $h_2 = 1$, we obtain $\mathcal{B}(r, h_1, 1) = \{3\mathbb{P}^{r-h_1+1}, \mathbb{P}^{r-h_1+2}\}$, hence $d(r, h_1, 1) = 2(r+h_1+1)$ and $g(r, h_1, 1) = 0$. For $s \geq 3$ and $h_s = 1$, we can suppose that $\mathbb{P}^{r+h_1} \cap \mathbb{P}^{2r-h_{(s-1)}+1} = \mathbb{P}^{r+h_1-h_{(s-1)}+1}$. Then the scroll degenerates into

1. $\mathcal{B}(r, h_1 - h_{(s-1)} + 1, \dots, h_{(s-2)} - h_{(s-1)} + 1, 1) \subset \mathbb{P}^{2r+1}$,
2. $\mathcal{B}(r-1, h_1, h_2-1, \dots, h_{(s-1)}-1, 1) \subset \mathbb{P}^{2r-1}$, with $\sigma(r-1)^2 \sigma(r+h_1-h_{(s-1)}) \sigma(r-h_1+h_2-2) \sigma(2r-h_2+h_3-2) \dots \sigma(2r-h_{(s-2)}+h_{(s-1)}-2)_{|2r-1}$ generators in common.

We set $\delta(r, h_1, \dots, h_{(s-2)}, 1) := \sigma(r)^2 \sigma(r+h_1) \sigma(r-h_1+h_2-1) \sigma(2r-h_2+h_3) \dots \sigma(2r-h_{(s-3)}+h_{(s-2)}) \sigma(2r-h_{(s-2)}+1)_{|2r+1}$. From Pieri's formula,

$$\delta(r, h_1, \dots, h_{(s-2)}, 1) = \sum_{k=0}^{h_{(s-2)}-1} \delta(r-k, h_1-k, h_2-2k, \dots, h_{(s-3)}-2k, 1)$$

Our assertion follows by induction on s because it is clear that $\delta(r, h_1, 1) = r-h_1+1$. Then

$$\begin{aligned} \bullet \quad d(r, h_1, \dots, h_{(s-1)}, 1) &= \sum_{k=0}^{h_{(s-1)}-1} d(r-k, h_1-h_{(s-1)}+1+k, h_2-h_{(s-1)}+1, \dots, h_{(s-2)}-h_{(s-1)}+1, 1), \\ \bullet \quad g(r, h_1, \dots, h_{(s-1)}, 1) &= \sum_{k=0}^{h_{(s-1)}-1} g(r-k, h_1-h_{(s-1)}+1+k, h_2-h_{(s-1)}+1, \dots, h_{(s-2)}-h_{(s-1)}+1, 1) + \sum_{k=0}^{h_{(s-1)}-2} \delta'(r-k-1, h_1-h_{(s-1)}+1+k, h_2-h_{(s-1)}+1, \dots, h_{(s-2)}-h_{(s-1)}+1, 1). \end{aligned}$$

By induction on $s \geq 2$, we obtain:

$$\begin{aligned}
\bullet \quad d(r, h_1, \dots, h_{(s-1)}, 1) &= 2 \sum_{k_1=0}^{h_{(s-1)}-1} \sum_{k_2=0}^{h_{(s-2)}-h_{(s-1)}} \dots \sum_{k_{(s-2)=0}^{h_2-h_3} (r-h_1+h_2-2 \sum_{i=1}^{s-2} k_i), \\
\bullet \quad g(r, h_1, \dots, h_{(s-1)}, 1) &= \sum_{k=0}^{h_{(s-1)}-2} \delta'(r-k-1, h_1-h_{(s-1)}+1+k, h_2-h_{(s-1)}+ \\
&\quad 1, \dots, h_{(s-2)}-h_{(s-1)}+1, 1) + \sum_{\alpha=2}^{s-2} \left[\sum_{k_1=0}^{h_{(s-1)}-1} \sum_{k_2=0}^{h_{(s-2)}-h_{(s-1)}} \dots \sum_{k_{(\alpha-1)=0}^{h_{(s-\alpha+1)}-h_{(s-\alpha+2)}} \right. \\
&\quad \left. \sum_{k_\alpha=0}^{h_{(s-\alpha)}-h_{(s-\alpha+1)}-1} \delta'(r-1-\sum_{i=1}^{\alpha} k_i, h_1-h_{(s-\alpha)}+1+\sum_{i=1}^{\alpha} k_i, h_2-h_{(s-\alpha)}+1, \dots, \right. \\
&\quad \left. h_{(s-\alpha-1)}-h_{(s-\alpha)}+1, 1) \right].
\end{aligned}$$

Finally, take $h_s \geq 2$. In $\mathcal{B}(r, h_1, \dots, h_s)$, suppose that $\mathbb{P}^{r+h_1} \cap \mathbb{P}^{2r-1} = \mathbb{P}^{r+h_1-1}$. Then the scroll degenerates into $\mathcal{B}(r, h_1-1, \dots, h_s-1) \subset \mathbb{P}^{2r+1}$ and $\mathcal{B}(r-1, h_1, h_2-1, \dots, h_s-1) \subset \mathbb{P}^{2r-1}$ with $\Delta_{\mathfrak{e} \not\sim 0}(r-1, h_1, h_2-1, \dots, h_s-1)$ generators in common. For this,

$$\begin{aligned}
\bullet \quad d(r, h_1, \dots, h_s) &= \sum_{k=0}^{h_s-1} \binom{h_s-1}{k} d(r-k, h_1-h_s+1+k, h_2-h_s+1, \dots, h_{(s-1)}-h_s+ \\
&\quad h_s+1, 1), \\
\bullet \quad g(r, h_1, \dots, h_s) &= \sum_{k=0}^{h_s-1} \binom{h_s-1}{k} g(r-k, h_1-h_s+1+k, h_2-h_s+1, \dots, h_{(s-1)}-h_s+ \\
&\quad 1, 1) + \sum_{k_1=0}^{h_s-2} \sum_{k_2=0}^{k_1} \binom{k_1}{k_2} \Delta'_{\mathfrak{e} \not\sim 0}(r-k_2-1, h_1-k_1+k_2, h_2-1-k_1, \dots, h_s-1-k_1). \blacksquare
\end{aligned}$$

In case $s = 2$, if $(r-h_1, r+h_1-h_2, 1^{(h_2-1)})$ is a partition of $2r-1$, then it is required that $2h_1 \leq h_2$. But the above formulas are true for all positive integers h_1, h_2 . This is possible because $\mathcal{B}(r, h_1, h_2)$ is base of an incidence scroll although $(r-h_1, r+h_1-h_2, 1^{(h_2-1)})$ is not a partition.

In the general case, all incidence scroll with $\mathfrak{e} \not\sim 0$ is one $R_g^d \subset \mathbb{P}^{2r-e+1+j}$ with $\{\mathbb{P}^{r-e}, \mathbb{P}^{r+j}\} \subset \mathcal{B}$ ($r := m-g+i_1$ and $j := i_2-i_1$). Moreover, there are a one-to-one correspondence between partitions of $2r+e+j+1$ such that $\lambda_1 \leq r-e$ and bases of such scrolls.

Lemma 3.9 *Let $\{h_1, \dots, h_s\}$ be a finite number of positive integers such that $0 \leq h_1 \leq r-e-1$, $1 \leq h_2 \leq r+j+h_1$, $1 \leq h_i \leq h_{(i-1)}-1$ for $3 \leq i \leq s$ and $h_s + \sum_{i=3}^s (h_{(i-1)}-h_i) = h_2$. Define $\Delta_{e \geq 1}(r, h_1, \dots, h_s) :=$*

$$\begin{cases} \sigma(r-e) \cdot \sigma(r+j) \cdot \sigma(r+j+h_1-1) \cdot \sigma(2r-e+j-1)^{(r+j+h_1-1)}, & s = 1 \\ \sigma(r-e) \cdot \sigma(r+j) \cdot \sigma(r+j+h_1-1) \cdot \sigma(r-e-h_1+h_2) \cdot \\ \sigma(2r-e+j-h_2+h_3) \cdot \dots \cdot \sigma(2r-e+j-h_{(s-1)}+h_s) \cdot \\ \sigma(2r-e+j-1)^{(h_s-1)}, & s \geq 2 \end{cases}$$

an intersection of Schubert cycles in $\mathbb{P}^{2r-e+j+1}$. Then:

$$\begin{aligned}
1. \Delta_{e \geq 1}(r, h) &= \sum_{k_1=0}^{r-e-h+1} \sum_{k_2=0}^{r-e-h+2-k_1} \cdots \sum_{k_h=0}^{r-e-\sum_{j=1}^{h-1} k_j} d_{e \geq 1}(r - \sum_{j=1}^h k_j, 0)_{(j-1)}; \\
2. \Delta_{e \geq 1}(r, h_1, 1) &= \begin{cases} r - e + 1 & \text{if } h_1 = 0 \\ r - e - h_1 + 2 & \text{if } h_1 > 0 \end{cases}; \\
3. \Delta_{e \geq 1}(r, h_1, \dots, h_s) &= \sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \sum_{k_3=0}^{h_{(s-2)}-h_s-2k_2} \cdots \sum_{k_{(s-1)}=0}^{h_2-h_s-2\sum_{j=2}^{s-2} k_j} \binom{h_s-1}{k_1} \\
\Delta_{e \geq 1}(r - \sum_{j=2}^{s-1} k_j, h_1 - h_s + 1 + k_1 - \sum_{j=2}^{s-1} k_j, 1), \quad s \geq 3.
\end{aligned}$$

The number $d_{e \geq 1}(r, 0)_j$ will be computed in Proposition 3.10.

Proof. If $s = 1$, then $\Delta(r, 0) = \sigma(r - e)\sigma(r + j - 1)^2\sigma(2r - e + j - 2)_{|2r-e+j}^{r+j-1} = d_{e \geq 1}(r, 0)_{j-1}$ (see Proposition 3.10). In general ($h \geq 1$), apply Pieri's formula to $\sigma(r + j + h - 1)$ and any $\sigma(2r - e + j - 1)$. Then, by induction on $r \geq e + h + 1$,

$$\Delta(r, h) = \Delta(r - 1, h) + \Delta(r, h - 1) = \sum_{k=0}^{r-e-h+1} \Delta(r - k, h - 1).$$

Use that $\Delta(e + h - 1, h) = \Delta(e + h - 1, h - 1)$.

For $s = 2$, $\Delta(r, 0, 1) = r - e + 1$ because is the degree of the directrix curve, which is contained in an $\mathbb{P}^{r-e+1}$, of $R_0^{2(r-e)+1}\mathbb{P}^{2(r-e)+2}$; and $\Delta(r, h_1 \geq 1, 1) = \sigma(r - e - h_1 + 1)_{|2(r-e-h_1+1)+1}^4 = r - e - h_1 + 2$.

The other cases are left to the reader because these are analogous of those of Lemma 3.3. $\blacksquare$

Proposition 3.10 *Each partition $(r - e - h, 1^{(r+j+h-1)})$ of $2r - e + j - 1$ such that $h \geq 0$ give rise to $R_{g_{e \geq 1}(r, h)}^{d_{e \geq 1}(r, h)} \subset \mathbb{P}^{2r-e+j+1}$ with base $\mathcal{B}_{e \geq 1}(r, h) = \{\mathbb{P}^{r-e}, \mathbb{P}^{r+j}, \mathbb{P}^{r+j+h}, (r + j + h - 1) \mathbb{P}^{2r-e+j-1}\}$ in general position, where*

$$\begin{aligned}
\bullet d_{e \geq 1}(r, h) &= \sum_{k_1=0}^{r-e-h} \sum_{k_2=0}^{r-e-h+1-k_1} \cdots \sum_{k_{(h-1)}=0}^{r-e-2-\sum_{i=1}^{h-2} k_i} \sum_{k_h=0}^{r-e-1-\sum_{i=1}^{h-1} k_i} \left[1 + \sum_{k_{(h+1)}=0}^{r-e-1-\sum_{i=1}^h k_i} \right. \\
&\quad \sum_{k_{(h+2)}=0}^{r-e-1-\sum_{i=1}^{h+1} k_i} \cdots \sum_{k_{(h+e+j-1)}=0}^{r-e-1-\sum_{i=1}^{h+e+j-2} k_i} d_{\mathfrak{e} \neq 0}(r - e - \sum_{i=1}^{h+e+j-1} k_i, 1) + \sum_{\alpha=0}^{j+e-2} \left[\sum_{k_{(h+1)}=0}^{r-e-1-\sum_{i=1}^h k_i} \cdots \right. \\
&\quad \left. \sum_{k_{(h+\alpha)}=0}^{r-e-1-\sum_{i=1}^{h+\alpha-1} k_i} (r - e - \sum_{i=1}^{h+\alpha} k_i) \right] \Big], \\
\bullet g_{e \geq 1}(r, h) &= \sum_{k_1=0}^{r-e-h} \sum_{k_2=0}^{r-e-h+1-k_1} \cdots \sum_{k_{(h-1)}=0}^{r-e-2-\sum_{i=1}^{h-2} k_i} \sum_{k_h=0}^{r-e-1-\sum_{i=1}^{h-1} k_i} \left[\sum_{k_{(h+1)}=0}^{r-e-2-\sum_{i=1}^h k_i} \cdots \right. \\
&\quad \sum_{k_{(h+e+j-1)}=0}^{r-e-2-\sum_{i=1}^{h+e+j-2} k_i} g_{\mathfrak{e} \neq 0}(r - e - \sum_{i=1}^{h+e+j-1} k_i, 1) + \sum_{\alpha=1}^{j+e-1} \left[\sum_{k_{(h+1)}=0}^{r-e-2-\sum_{i=1}^h k_i} \cdots \right.
\end{aligned}$$

$$\begin{aligned}
& \sum_{k_{(h+\alpha)}=0}^{r-e-2-\sum_{i=1}^{h+\alpha-1} k_i} \beta(r-1-\sum_{i=1}^{h+\alpha} k_i, j-\alpha) \Big] + \sum_{k=0}^{r-e-h-1} \Delta'_{e \geq 1}(r-k-1, h) + \sum_{\alpha=2}^{h-1} \left[\sum_{k_1=0}^{r-e-h} \right. \\
& \sum_{k_2=0}^{r-e-h+1-k} \cdots \sum_{k_{(\alpha-1)}=0}^{r-e-h+\alpha-2-\sum_{i=1}^{\alpha-2} k_i} \sum_{k_\alpha=0}^{r-e-h+\alpha-2-\sum_{i=1}^{\alpha-1} k_i} \Delta'_{e \geq 1}(r-1-\sum_{i=1}^{\alpha} k_i, h-\alpha+1) \Big],
\end{aligned}$$

with $\beta(r, j) = d(r, 0)_j - 1$.

Proof. We proceed by induction on $h \geq 0$. For $h = 0$, let us the temporary notation of $(r, 0)_j$ for $(r, 0)$. It is clear that $r+h+j > 1$. In other case, we have $\mathcal{B} = \{3\mathbb{P}^1\} \subset \mathbb{P}^3$ but $e \not\sim 0$. In $\mathcal{B}(r, 0)_j$, suppose that $\mathbb{P}^{r-e} \cap \mathbb{P}^{2r-e+j-1} = \mathbb{P}^{r-e-1}$. Then the scroll breaks up into $\mathcal{B}(r, 0)_{j-1} \subset \mathbb{P}^{2r-e+j}$ and $\mathcal{B}(r-1, 0)_j \subset \mathbb{P}^{2r-e+j-1}$ with $\sigma(r-e-1)\sigma(r+j-1)^2\sigma(2r-e+j-2)^{r+j-2} = \sigma(r-e-1)\sigma(r+j-2)^2\sigma(2r-e+j-4)^{r+j-2} = d(r-1, 0)_{j-1}$ generators in common. By induction on $r \geq e+1$, we find that

$$\begin{aligned}
& \bullet d(r, 0)_j = 1 + \sum_{k=0}^{r-e-1} d(r-k, 0)_{j-1}, \\
& \bullet g(r, 0)_j = \sum_{k=0}^{r-e-2} g(r-k, 0)_{j-1} + \sum_{k=0}^{r-e-2} [d(r-1-k, 0)_{j-1} - 1].
\end{aligned}$$

When $j = -e+1$, it is easy to check that $d(r, 0)_{(-e+1)} = 1 + d\mathfrak{e}_{\not\sim 0}(r-e, 1)$ and $g(r, 0)_{(-e+1)} = g\mathfrak{e}_{\not\sim 0}(r-e, 1)$. Hence, by induction on $j \geq -e+1$,

$$\begin{aligned}
& \bullet d(r, 0)_j = 1 + \sum_{k_1=0}^{r-e-1} \sum_{k_2=0}^{r-e-1-k_1} \cdots \sum_{k_{(j+e-1)}=0}^{r-e-1-\sum_{i=1}^{j+e-2} k_i} d\mathfrak{e}_{\not\sim 0}(r-e-\sum_{i=1}^{j+e-1} k_i, 1) + \\
& \sum_{\alpha=0}^{j+e-2} \left[\sum_{k_1=0}^{r-e-1} \sum_{k_2=0}^{r-e-1-k_1} \cdots \sum_{k_\alpha=0}^{r-e-1-\sum_{i=1}^{\alpha-1} k_i} (r-e-\sum_{i=1}^{\alpha} k_i) \right], \\
& \bullet g(r, 0)_j = \sum_{k_1=0}^{r-e-2} \sum_{k_2=0}^{r-e-2-k_1} \cdots \sum_{k_{(j+e-1)}=0}^{r-e-2-\sum_{i=1}^{j+e-2} k_i} g\mathfrak{e}_{\not\sim 0}(r-e-\sum_{i=1}^{j+e-1} k_i, 1) + \sum_{\alpha=1}^{j+e-1} \\
& \left[\sum_{k_1=0}^{r-e-2} \sum_{k_2=0}^{r-e-2-k_1} \cdots \sum_{k_\alpha=0}^{r-e-2-\sum_{i=1}^{\alpha-1} k_i} \beta(r-1-\sum_{i=1}^{\alpha} k_i, j-\alpha) \right].
\end{aligned}$$

Assuming the formulas hold for $h-1 \geq 0$, we will prove them for h . In $\mathcal{B}(r, h)$, suppose that $\mathbb{P}^{r+j+h} \cap \mathbb{P}^{2r-e+j-1} = \mathbb{P}^{r+j+h-1}$. Then R degenerates into $\mathcal{B}(r, h-1) \subset \mathbb{P}^{2r-e+j+1}$ and $\mathcal{B}(r-1, h) \subset \mathbb{P}^{2r-e+j-1}$ with $\Delta_{e \geq 1}(r-1, h)$ generators in common. By induction on $r \geq e+h+1$, we find that

$$\bullet d(r, h) = \sum_{k=0}^{r-e-h} d(r-k, h-1) = \sum_{k_1=0}^{r-e-h} \sum_{k_2=0}^{r-e-h+1-k_1} \cdots \sum_{k_h=0}^{r-e-1-\sum_{i=1}^{h-1} k_i} d(r-\sum_{i=1}^h k_i, 0),$$

$$\begin{aligned}
\bullet \quad g(r, h) &= \sum_{k=0}^{r-e-h} g(r-k, h-1) + \sum_{k=0}^{r-e-h-1} \Delta'(r-k-1, h) = \sum_{k_1=0}^{r-e-h} \sum_{k_2=0}^{r-e-h-1-k_1} \dots \\
&\quad \sum_{k_h=0}^{r-e-1-\sum_{i=1}^{h-1} k_i} g(r - \sum_{i=1}^h k_i, 0) + \sum_{\alpha=2}^{h-1} \left[\sum_{k_1=0}^{r-e-h} \sum_{k_2=0}^{r-e-h-1-k_1} \dots \sum_{k_{(\alpha-1)}=0}^{r-e-h+\alpha-2-\sum_{i=1}^{\alpha-2} k_i} \right. \\
&\quad \left. \sum_{k_\alpha=0}^{r-e-h+\alpha-2-\sum_{i=1}^{\alpha-1} k_i} \Delta'(r-1-\sum_{i=1}^{\alpha} k_i, h-\alpha+1) \right] + \sum_{k=0}^{r-e-h-1} \Delta'(r-k-1, h);
\end{aligned}$$

The last equalities can be proved by induction on $h \geq 0$. $\blacksquare$

Remark 3.11 For $\mathcal{B}_{e \geq 1}(r, h_1, h_2) = \{\mathbb{P}^{r-e}, \mathbb{P}^{r+j}, \mathbb{P}^{r+j+h_1}, \mathbb{P}^{r-e-h_1+h_2}, (h_2-1)\mathbb{P}^{2r-e+j-1}\}$ in general position, it follows easily that

$$d_{e \geq 1}(r, h_1, 1) = \begin{cases} 2(r-e)+1 & \text{if } h_1 = 0 \\ 2(r-e-h_1)+2 & \text{if } h_1 > 0 \end{cases} \quad ; \quad g_{e \geq 1}(r, h_1, 1) = 0.$$

If $h_2 = 1$, then $e+j+2h_1 \leq 1$ but $e+j > 0$, i.e., $h_1 = 0$. For this $d(r, 0, 1) = 2(r-e)+1$ and $g(r, 0, 1) = 0$. In case $h_1 \geq 1$, we obtain $\mathcal{B}(r, h_1, 1) = \{3\mathbb{P}^{r-e-h_1+1}, \mathbb{P}^{r-e-h_1+2}\} \subset \mathbb{P}^{2(r-e-h_1+1)+1}$, i.e., $d(r, h_1, 1) = 2(r-e-h_1+1)$ and $g(r, h_1, 1) = 0$. This result will be needed in the following theorem which may be proved in much the same way as Theorem 3.8

Theorem 3.12 For $s \geq 2$, each partition $(r-e-h_1, r+j+h_1-h_2, h_2-h_3, \dots, h_{(s-1)}-h_s, 1^{(h_s-1)})$ of $2r-e+j-1$ such that $h_1 \geq 0$ corresponds to the scroll $R_{g_{e \geq 1}(r, h_1, \dots, h_s)}^{d_{e \geq 1}(r, h_1, \dots, h_s)} \subset \mathbb{P}^{2r-e+j+1}$ with base $\mathcal{B}_{e \geq 1}(r, h_1, \dots, h_s) = \{\mathbb{P}^{r-e}, \mathbb{P}^{r+j}, \mathbb{P}^{r+j+h_1}, \mathbb{P}^{r-e-h_1+h_2}, \mathbb{P}^{2r-e+j-h_2+h_3}, \dots, \mathbb{P}^{2r-e+j-h_{(s-1)}+h_s}, (h_s-1)\mathbb{P}^{2r-e+j-1}\}$ in general position, where

$$\begin{aligned}
\bullet \quad d_{e \geq 1}(r, h_1, \dots, h_s) &= \sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \dots \sum_{k_{(s-1)}=0}^{h_2-h_3} \binom{h_s-1}{k_1} d(r - \sum_{i=1}^{s-1} k_i, h_1-h_2+1 + \sum_{i=1}^{s-1} k_i, 1), \\
\bullet \quad g_{e \geq 1}(r, h_1, \dots, h_s) &= \sum_{\alpha=1}^{s-2} \left[\sum_{k_1=0}^{h_s-1} \sum_{k_2=0}^{h_{(s-1)}-h_s} \dots \sum_{k_\alpha=0}^{h_{(s-\alpha+1)}-h_{(s-\alpha+2)}} \sum_{k_{(\alpha+1)}=0}^{h_{(s-\alpha)}-h_{(s-\alpha+1)}-1} \right. \\
&\quad \left. \binom{h_s-1}{k_1} [\delta(r-1-\sum_{i=1}^{\alpha+1} k_i, h_1-h_{(s-\alpha)}+1 + \sum_{i=1}^{\alpha+1} k_i, h_2-h_{(s-\alpha)}+1, h_{(s-\alpha-1)}-h_{(s-\alpha)}+1, 1) - 1] \right] \\
&\quad + \sum_{k_1=0}^{h_s-2} \sum_{k_2=0}^{k_1} \binom{k_1}{k_2} \Delta'_{e \geq 1}(r-k_2-1, h_1-k_1+k_2, h_2-1-k_1, \dots, h_s-1-k_1),
\end{aligned}$$

letting $\beta'(r, h_1, \dots, h_{(s-2)}, 1) = \delta'(r-e, h_1, \dots, h_{(s-2)}, 1)$. $\blacksquare$

Remark 3.13 In general, $\mathcal{B}_{e \geq 1}(r, h_1, \dots, h_s) \subset \mathbb{P}^{2r-e+j+1}$ is not base of an incidence scroll which corresponds to an abstract model with $\deg(-\mathfrak{E}) = e$.

For example, for $s = 1$, $h_1 = 0$ and $r = e + 1$, we know that $\mathcal{B}(e + 1, 0)$ generates $R_0^{e+j+2} \subset \mathbb{P}^{e+j+3}$. It is the immersion of the rational ruled surface $X_{(e+j)} = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-e-j))$ by the very ample divisor $H \sim C_o + (e+j)f$, i.e., $\deg(-\mathfrak{E}) = e+j \geq 1$. For this reason, if we talk about $\mathbb{P}^{2r-e+j+1}$, then we need only consider the numbers r, e and j such that $r+j > r-e \geq 1$. So we obtain a decomposable incidence scroll with $\deg(-\mathfrak{E}) \geq 1$.

Theorem 3.14 *Let $\pi : X = \mathbb{P}(\mathcal{E}) \rightarrow C$ be a decomposable ruled surface over a curve C of genus g , such that $h^0(\mathcal{O}_X(C_0 - \mathfrak{E}f)) = 1$. Let $\tilde{X}_x = \mathbb{P}(\tilde{\mathcal{E}})$ be the elementary transform of X at x where $x \notin C_o$ and $x \notin C_1$. Then, every projective model of X as an incidence scroll $R_g^d \subset \mathbb{P}^n$ with base $\mathcal{B} = \{\mathbb{P}^{n_1}, \mathbb{P}^{n_2}, \dots, \mathbb{P}^{n_r}\}$ in general position, gives rise to an projective model of $\tilde{X}_x$, which is an incidence scroll $R_g^{d-1} \subset \mathbb{P}^{n-1}$ with base $\tilde{\mathcal{B}} = \{\mathbb{P}^{n_1}, \mathbb{P}^{n_2}, \mathbb{P}^{n_3-1}, \dots, \mathbb{P}^{n_r-1}\}$ in general position.*

Furthermore, these are all the indecomposable incidence scrolls of $\mathbb{P}^{n-1}$ with $n_1 + n_2 = n - 1$.

Proof. Let $\Phi_{|H|} : X \rightarrow R_g^d \subset \mathbb{P}^n$ be the immersion induced by $H \sim C_o + \mathfrak{b}f$ on X . Let $P = \Phi_{|H|}(x)$ with $y = \pi(x)$. Then the elementary transform of X at x corresponds to the projection of $R_g^d \subset \mathbb{P}^n$ from P . Hence

$$\begin{array}{ccc} X = \mathbb{P}(\mathcal{E}) & \xrightarrow{\phi_{|C_o + \mathfrak{b}f|}} & R_g^d \subset \mathbb{P}^n \\ \xi_x^{-1} \downarrow & & \downarrow \pi_P \\ \tilde{X}_x = \mathbb{P}(\tilde{\mathcal{E}}) & \xrightarrow{\phi_{|\xi^*(C_o) + (\mathfrak{b}-y)f|}} & R_g^{d-1} \subset \mathbb{P}^{n-1} \end{array}$$

Moreover, project $R_g^d \subset \mathbb{P}^n$ from P , such that $P \notin C_g^{m-e} \subset \mathbb{P}^{n_1}$ and $P \notin C_g^m \subset \mathbb{P}^{n_2}$ for any C_g^m , is equivalent to join $\mathbb{P}^{n_1}$ and $\mathbb{P}^{n_2}$. Then, we obtain an incidence scroll $R_g^{d-1} \subset \mathbb{P}^{n-1}$ with base $\tilde{\mathcal{B}}$ which is indecomposable because $n_2 < n_3$, or equivalently $h^0(\mathcal{O}_X(C_0 - \mathfrak{E}f)) = 1$. ■

In this way, we can now compute the degree and genus of these particular incidence scroll, using the above results for decomposable incidence scrolls and Proposition 1.3.

References

- [1] CID-MUÑOZ, R.-PEDREIRA, M. *Classification of Incidence Scrolls(I)*
Manuscripta Mathematica 105, pp. 125-138 (2001).
- [2] CID-MUÑOZ, R.-PEDREIRA, M. *Another Classification of Incidence Scrolls*
Accepted in Archives of Mathematics, preprint: AG/0201193.

- [3] FULTON, W. *Intersection Theory*.
Ergebnisse der Mathematik und ihrer Grenzgebiete (3), Springer-Verlag,
Berlin (1984).
- [4] FULTON, W. *Young Tableaux with Applications to Representation Theory and Geometry*.
London Mathematical Society Student texts 35 (1997).
- [5] GRIFFITHS, P.-HARRIS, J. *Principles of Algebraic Geometry*.
Wiley-Interscience, New York (1978), reprinted (1994).
- [6] HARTSHORNE, R. *Algebraic Geometry*.
Graduate Texts in Mathematics 52. Springer-Verlag, New York. 1977.
- [7] HODGE, W.V.D.-PEDOE, D. *Methods of Algebraic Geometry II*.
Cambridge University Press (1952), reprinted (1968).
- [8] KLEIMAN, S. L.-LAKSOV, D. *Schubert calculus*.
Amer. Math. Monthly 79, pp. 1061-1082 (1972).
- [9] KLEIMAN, S. L. *The transversality of a general translate*.
Compositio Mathematica, volume 28, pp. 287-297 (1974).
- [10] SOTTILE, F. *Explicit Enumerative Geometry for the Real Grassmannian of Lines in Projective Space*.
Duke Mathematical Journal, volume 87, pp. 59-85 (1997).